AD					

Award Number: W81XWH-05-1-0137

TITLE: A Role for TACI in Prostate Neoplasia

PRINCIPAL INVESTIGATOR: Gotz-Ulrich von Bulow, Ph.D.

CONTRACTING ORGANIZATION: Indiana University

Indianapolis, IN 46202-5167

REPORT DATE: January 2008

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

**Distribution Unlimited** 

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

# REPORT DOCUMENTATION PAGE Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS. 1. REPORT DATE (DD-MM-YYYYY) 2. REPORT TYPE 3. DATES COVERED (From - To)

1. REPORT DATE (DI	LEASE DO NOT RETURN YOU						
	*	2. REPORT TYPE			ATES COVERED (From - To)		
01-01-2008		Final			Dec 2005 – 14 Dec 2007		
4. TITLE AND SUBTIT	LE			5a.	CONTRACT NUMBER		
A Role for TACI i	n Prostate Neoplas	sia			GRANT NUMBER		
					1XWH-05-1-0137		
				5c.	PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d.	PROJECT NUMBER		
Gotz-Ulrich von Br	ulow, Ph.D.			5e. `	TASK NUMBER		
E-Mail: gvonbulo	@iupui.edu			5f. \	WORK UNIT NUMBER		
7. PERFORMING ORG	SANIZATION NAME(S)	AND ADDRESS(ES)		8. P	ERFORMING ORGANIZATION REPORT		
	` '	` ,		N	UMBER		
Indiana University							
Indianapolis, IN 4	6202-5167						
, ,							
a SPONSOPING / MC	MITODING AGENCY I	NAME(S) AND ADDRES	S/ES)	10	SPONSOR/MONITOR'S ACRONYM(S)		
	I Research and Ma		3(E3)	10.	SPONSOR/MONTOR S ACRONTM(S)		
•		ilenei Command					
Fort Detrick, Mary	and 21/02-5012			44			
					SPONSOR/MONITOR'S REPORT		
					NUMBER(S)		
	VAILABILITY STATE						
Approved for Publ	ic Release; Distribι	ution Unlimited					
13. SUPPLEMENTAR	Y NOTES – Original c	contains colored plates	: ALL DTIC reproduc	ctions will be in	black and white.		
10. OOI I ELINENTAN	THOTEG Original c	onams colored plates	. TIEE DITE reproduc	ctions will be in	orack and write.		
14. ABSTRACT							
14. ABOTTAGT							
	oposed that TNF- family	growth factors, BAFF ar	nd APRIL play a role in pr	rostrate cancer. C	our hypothesis proposed that APRIL provides		
This research study pro	pposed that TNF- family normal prostate epithel	r growth factors, BAFF ar ial cells by means of an ι	nd APRIL play a role in pr unknown receptor. We po	rostrate cancer. Costulated that TAC	our hypothesis proposed that APRIL provides		
This research study pro a proliferative signal to to regulate APRIL-med	normal prostate epithel liated proliferation of cel	lial cells by means of an u lls by inducing apoptosis	unknown receptor. We po and thereby maintaining	ostulated that TAC the normal glandu	I is an antagonist receptor which functions lar structure. The loss of TACI expression in		
This research study pro a proliferative signal to to regulate APRIL-med prostate cells therefore	normal prostate epithel liated proliferation of cel conceivably results in	lial cells by means of an u lls by inducing apoptosis an imbalance in homeost	unknown receptor. We po and thereby maintaining asis resulting in the aber	ostulated that TAC the normal glandu rant accumulation	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to		
This research study pro- a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy	normal prostate epithel liated proliferation of cel conceivably results in a pothesis anticipated tha	lial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w	Inknown receptor. We po and thereby maintaining asis resulting in the aber ould enhance cell growth	ostulated that TAC the normal glandu rant accumulation n whereas the add	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to tion of TACI-Ig would either reduce cell		
This research study pro- a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apop	normal prostate epithel liated proliferation of cel conceivably results in a pothesis anticipated tha tosis. This did not happ	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	unknown receptor. We po and thereby maintaining asis resulting in the aber ould enhance cell growth by significant changes in	ostulated that TAC the normal glandu rant accumulation n whereas the add the relative numbe	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to ition of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by		
This research study pro- a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apopt propidium iodide flow of	normal prostate epithel liated proliferation of cel conceivably results in a pothesis anticipated tha tosis. This did not happ	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	unknown receptor. We po and thereby maintaining asis resulting in the aber ould enhance cell growth by significant changes in	ostulated that TAC the normal glandu rant accumulation n whereas the add the relative numbe	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to tion of TACI-Ig would either reduce cell		
This research study pro- a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apop	normal prostate epithel liated proliferation of cel conceivably results in a pothesis anticipated tha tosis. This did not happ	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	unknown receptor. We po and thereby maintaining asis resulting in the aber ould enhance cell growth by significant changes in	ostulated that TAC the normal glandu rant accumulation n whereas the add the relative numbe	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to ition of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by		
This research study pro- a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apopt propidium iodide flow of	normal prostate epithel liated proliferation of cel conceivably results in a pothesis anticipated tha tosis. This did not happ	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	unknown receptor. We po and thereby maintaining asis resulting in the aber ould enhance cell growth by significant changes in	ostulated that TAC the normal glandu rant accumulation n whereas the add the relative numbe	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to ition of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by		
This research study pro- a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apopt propidium iodide flow of	normal prostate epithel liated proliferation of cel conceivably results in a pothesis anticipated tha tosis. This did not happ	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	unknown receptor. We po and thereby maintaining asis resulting in the aber ould enhance cell growth by significant changes in	ostulated that TAC the normal glandu rant accumulation n whereas the add the relative numbe	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to ition of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by		
This research study pro- a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apopt propidium iodide flow of	normal prostate epithel liated proliferation of cel conceivably results in a pothesis anticipated tha tosis. This did not happ	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	unknown receptor. We po and thereby maintaining asis resulting in the aber ould enhance cell growth by significant changes in	ostulated that TAC the normal glandu rant accumulation n whereas the add the relative numbe	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to ition of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by		
This research study pro- a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apopt propidium iodide flow of	normal prostate epithel liated proliferation of cel conceivably results in a pothesis anticipated tha tosis. This did not happ	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	unknown receptor. We po and thereby maintaining asis resulting in the aber ould enhance cell growth by significant changes in	ostulated that TAC the normal glandu rant accumulation n whereas the add the relative numbe	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to ition of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by		
This research study pro- a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apopt propidium iodide flow of	normal prostate epithel liated proliferation of cel conceivably results in a pothesis anticipated tha tosis. This did not happ	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	unknown receptor. We po and thereby maintaining asis resulting in the aber ould enhance cell growth by significant changes in	ostulated that TAC the normal glandu rant accumulation n whereas the add the relative numbe	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to ition of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by		
This research study pro a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apop propidium iodide flow of Indiana University	normal prostate epithel liated proliferation of cel e conceivably results in a pothesis anticipated tha losis. This did not happ cytometry. Because of the	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	unknown receptor. We po and thereby maintaining asis resulting in the aber ould enhance cell growth by significant changes in	ostulated that TAC the normal glandu rant accumulation n whereas the add the relative numbe	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to ition of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by		
This research study pro- a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apopt propidium iodide flow of	normal prostate epithel liated proliferation of cel e conceivably results in a pothesis anticipated tha losis. This did not happ cytometry. Because of the	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	unknown receptor. We po and thereby maintaining asis resulting in the aber ould enhance cell growth by significant changes in	ostulated that TAC the normal glandu rant accumulation n whereas the add the relative numbe	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to ition of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by		
This research study program a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apopy propidium iodide flow of Indiana University	normal prostate epithel liated proliferation of cel e conceivably results in a pothesis anticipated tha losis. This did not happ cytometry. Because of the	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	unknown receptor. We po and thereby maintaining asis resulting in the aber ould enhance cell growth by significant changes in	ostulated that TAC the normal glandu rant accumulation n whereas the add the relative numbe	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to ition of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by		
This research study program a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apopy propidium iodide flow of Indiana University	normal prostate epithel liated proliferation of cel e conceivably results in a pothesis anticipated tha losis. This did not happ bytometry. Because of the cytometry. Because of the	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	anknown receptor. We pound thereby maintaining asis resulting in the aberrould enhance cell growth by significant changes in rudy was terminated early	ostulated that TAC the normal glandurant accumulation whereas the add the relative number and the grant relative numbers.	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to tion of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by inquished. Dr. VonBulow no longer works for		
This research study program a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apopy propidium iodide flow of Indiana University  15. SUBJECT TERMS No subject terms provi	normal prostate epithel liated proliferation of cel e conceivably results in a pothesis anticipated tha losis. This did not happ bytometry. Because of the cytometry. Because of the	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	unknown receptor. We po and thereby maintaining asis resulting in the aber ould enhance cell growth ny significant changes in udy was terminated early	estulated that TAC the normal glandurant accumulation whereas the add the relative numbery and the grant rel	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to tion of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by inquished. Dr. VonBulow no longer works for		
This research study program a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apopy propidium iodide flow of Indiana University  15. SUBJECT TERMS No subject terms provi	normal prostate epithel liated proliferation of cel e conceivably results in a pothesis anticipated tha losis. This did not happ bytometry. Because of the cytometry. Because of the	ial cells by means of an u lls by inducing apoptosis an imbalance in homeost t the addition of APRIL w en and we did not see ar	anknown receptor. We pound thereby maintaining asis resulting in the aberrould enhance cell growth by significant changes in rudy was terminated early	ostulated that TAC the normal glandurant accumulation whereas the add the relative number and the grant relative numbers.	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to tion of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by inquished. Dr. VonBulow no longer works for		
This research study program a proliferative signal to to regulate APRIL-med prostate cells therefore transformation. Our hy growth or induce apopt propidium iodide flow of Indiana University  15. SUBJECT TERMS No subject terms provi	normal prostate epithel liated proliferation of cel e conceivably results in a pothesis anticipated tha losis. This did not happ cytometry. Because of the ded.	ial cells by means of an ulls by inducing apoptosis an imbalance in homeost it the addition of APRIL wen and we did not see are negative results, the st	anknown receptor. We pound thereby maintaining asis resulting in the aberrould enhance cell growth by significant changes in rudy was terminated early	ostulated that TAC the normal glandurant accumulation whereas the add the relative number and the grant relative from the grant from th	I is an antagonist receptor which functions lar structure. The loss of TACI expression in of cells which become susceptible to ition of TACI-Ig would either reduce cell ers of hyperdiploid cells as determined by inquished. Dr. VonBulow no longer works for USAMRMC		

# Table of Contents

Introduction	4
Body	4
Key Research Accomplishments	4
Reportable Outcomes	4
Conclusions	5
References	5
Appendices	5

#### Introduction

The goal of this grant was to gain insight into the molecular basis of prostate cancer. Preliminary evidence suggested that the taci gene is expressed in normal prostate tissues, but not in prostate tumor cells. We had proposed the APRIL provides a proliferative signal to normal prostate epithelial cells by means of an unknown receptor.

# Body

To determine the role of APRIL and TACI in prostate tumor growth, we (a)cultured LNCaP, PC3 and DU145 prostate cancer cell lines and titrated the effects of recombinant APRIL and TACL-Ig on cell growth as determined by MTT assay, DNA content of cells and Annexin V binding assays; (b) we did not determine the effect of systemic TACI-Ig administration on the growth of LNCaP cells in nude mice because of the negative results obtained in (a); (c) To develop LNCaP clones with tetracycline-inducible TACI expression (Tet-TACI) and determine the effects of TACI activation on cell growth and apoptosis as determined by DNA content and Annexin V binding assay, we utilized the commercial "Tet-on" system from BD-Clontech and were able to derive several inducible TACI-expressing clones for the LNCaP and CWR22cell lines. In the latter (c),data was repeated at least twice with three inducible clones from each cell line. These experiments were also repeated in conditions of reduced serum.

## **Key Research Accomplishments**

Unfortunately, we have no major research accomplishments to list.

#### **Reportable Outcomes**

(a) According to our hypothesis we anticipated that the addition of recombinant APRil would enhance cell growth whereas the addition of TACi-IL would either reduce cell growth or induce apoptosis. This did not happen and for all four cell lines tested we did not see any significant changes in relative cell number (ad determined by MTT reduction), nor did we see any significant changes in the relative numbers of hyperdiploid cells as determined by propidium iodide flow cytometry. These experiments were repeated several times with similar results. To overcome the possibility that the cells were already maximally stimulated in the presence off fetal calf serum, we repeated the MTT experiments in either serum-free conditions, or in medium with 1% fetal calf serum. Although the MTT values were reduced overall, there was no significant difference observed when titrating in either APRIL or TACI-Ig. We included the titration with BAFF since it is also a TACI ligand, but similar negative results were obtained with this cytokine. When performing the apoptosis assays with propidium iodide and annexin V staining, we encountered a pitfall in that we were not able to separate the cells from the monolayer into individual cells for flow cytometry using EDTA or trypsin without affecting the viability of the cells (and thereby propidium iodide staining). Overall these types of experiments were not very reproducible and were therefore abandoned as unreliable.

(b) Due to the negative results obtained in (a) and the large amount of resources required, we decided not to perform the experiments to determine the effect of systemic TACI-Ig administration on the growth of LNCaP cells in nude mice.

(c) Unfortunately no relative change of growth characteristics was observed in the presence of doxcycyclin when compared with the parental cell lines. Additionally, induction of TACI expression coupled with treatment with APRIL, BAFF, or TACI-IG also did not manifest any significant differences in cell growth as determined by MTT assay.

(d) Due to the negative results obtained in (c) we did not perform experiments to determine the effect of docycycline induction of TACI expression of Tet-TACI LNCaP cells inplanted in nude mice.

### **Conclusions**

The experiments conducted in this research netted essentially negative results and thus the research project was stopped early and the grant relinquished.

**References: None** 

**Appendices: None**